

Characterization of Balanced Coherent Configurations

Mitsugu Hirasaka^{1,*}, Reza Sharafadini^{1,2,*}

*Department of Mathematics, College of Natural Sciences, Pusan National University,
Busan 609-735, South Korea.*

Abstract

Let G be a group acting on a finite set Ω . Then G acts on $\Omega \times \Omega$ by its entry-wise action and its orbits form the basis relations of a coherent configuration (or shortly scheme). Our concern is to consider what follows from the assumption that the number of orbits of G on $\Omega_i \times \Omega_j$ is constant whenever Ω_i and Ω_j are orbits of G on Ω . One can conclude from the assumption that the actions of G on Ω_i 's have the same permutation character and are not necessarily equivalent. From this viewpoint one may ask how many inequivalent actions of a given group with the same permutation character there exist. In this article we will approach to this question by a purely combinatorial method in terms of schemes and investigate the following topics: (i) balanced schemes and their central primitive idempotents, (ii) characterization of reduced balanced schemes.

Keywords: Coherent configurations, (m, n, r) -schemes, Balanced schemes, Central primitive idempotents.

1. Introduction

Let G be a group acting on a finite set Ω with its orbits $\Omega_1, \dots, \Omega_n$ and its permutation character $\pi = \sum_{i=1}^n \pi_i$ where $\pi_i(g) := |\{\alpha \in \Omega_i \mid \alpha^g = \alpha\}|$ for $g \in G$. One may think what happens if $\pi_i = \pi_j$ for all $1 \leq i, j \leq n$ and

*Corresponding Author

Email addresses: hirasaka@pusan.ac.kr (Mitsugu Hirasaka),
sharafadini@pusan.ac.kr (Reza Sharafadini)

¹Supported by the Korean Research Foundation grant (KRF) No: 2009-0070854.

²Supported by BK21 Dynamic Math Center, Department of Mathematics at Pusan National University.

can say that the number of orbits of G on $\Omega_i \times \Omega_j$ by its entry-wise action is constant for all $1 \leq i, j \leq n$, which motivate us to define the following concepts whose terminology is due to [3].

Definition 1.1. Let V be a finite set and $\mathcal{R}$ a set of nonempty binary relations on V . The pair $\mathcal{C} = (V, \mathcal{R})$ is called a *coherent configuration* (for short *scheme*) on V if the following conditions hold:

- (C1) $\mathcal{R}$ forms a partition of the set $V \times V$.
- (C2) $\Delta_V := \{(v, v) \mid v \in V\}$ is a union of certain relations from $\mathcal{R}$.
- (C3) For every $R \in \mathcal{R}$, $R^t := \{(v, u) \mid (u, v) \in R\} \in \mathcal{R}$.
- (C4) For every $R, S, T \in \mathcal{R}$, the size of $\{w \in V \mid (u, w) \in R, (w, v) \in S\}$ does not depend on the choice of $(u, v) \in T$ and is denoted by c_{RS}^T .

We say that the elements of V are *points* and those of $\mathcal{R}$ are *basis relations*.

Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme and $\emptyset \neq X \subseteq V$. We say that X is a *fiber* of $\mathcal{C}$ if $\Delta_X = \{(x, x) \mid x \in X\} \in \mathcal{R}$. We denote by $\text{Fib}(\mathcal{C})$ the set of all fibers of $\mathcal{C}$.

Definition 1.2. Let m, n and r be positive integers. We say that a scheme $\mathcal{C}$ is an (m, n, r) -*scheme* if the following conditions hold:

- (i) $|\{R \in \mathcal{R} \mid R \subseteq X \times Y\}| = r$ for all $X, Y \in \text{Fib}(\mathcal{C})$.
- (ii) $|X| = m$ for all $X \in \text{Fib}(\mathcal{C})$.
- (iii) $|\text{Fib}(\mathcal{C})| = n$.

A scheme $\mathcal{C}$ is called *r-balanced* if (i) holds, and *balanced* if it is *r-balanced* for some r . In Section 3 we will show that (i) implies (ii).

Let us return to the topic in the first paragraph. Note that the orbits of G on $\Omega \times \Omega$ form the basis relations of a scheme called the *2-orbit scheme* of G on Ω and its fibers are $\Omega_1, \dots, \Omega_n$. Furthermore, if $\pi_i = \pi_j$ for all $1 \leq i, j \leq n$, then the 2-orbit scheme of G on Ω is balanced.

We denote by $\mathcal{P}(\mathcal{C})$ the set of all central primitive idempotents of the adjacency algebra of $\mathcal{C}$ (see Section 2 for details). The following theorem shows a characterization of balanced schemes in terms of their central primitive idempotents.

Theorem 1.1. *Let $\mathcal{C}$ be a scheme. Then $\mathcal{C}$ is balanced if and only if for each $X \in \text{Fib}(\mathcal{C})$ the mapping $\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C}_X)$ ($P \mapsto P_X$) is bijective with $n_P = |\text{Fib}(\mathcal{C})|n_{P_X}$.*

One may conclude that $|\mathcal{P}(\mathcal{C})| = r$ if $\mathcal{C}$ is r -balanced and $r \leq 5$ (see Corollary 3.1). The following theorem deals with the converse argument for $r = 1, 2$.

Theorem 1.2. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. Then the following hold:*

- (i) $|\mathcal{P}(\mathcal{C})| = 1$ if and only if $\mathcal{C}$ is 1-balanced.
- (ii) $|\mathcal{P}(\mathcal{C})| = 2$ if and only if $\mathcal{C} = \mathcal{C}_1 \boxplus \mathcal{C}_2$ where $\mathcal{C}_i$ is i -balanced.

We have the following constructions of balanced schemes (see Sections 3, 4 for the details):

- (i) Let U be a union of fibers of $\mathcal{C}$. Then the restriction of $\mathcal{C}$ to U is r -balanced if $\mathcal{C}$ is r -balanced.
- (ii) If $\mathcal{C}_i$ ($i = 1, 2$) is an (m_i, n_i, r_i) -scheme, then $\mathcal{C}_1 \otimes \mathcal{C}_2$ is an $(m_1 m_2, n_1 n_2, r_1 r_2)$ -scheme.

We say that a balanced scheme $\mathcal{C}$ is *reduced* if there exist no $X, Y \in \text{Fib}(\mathcal{C})$ such that $\mathcal{C}_{X \cup Y} \simeq \mathcal{C}_X \otimes \mathcal{T}_2$ where $\mathcal{T}_2$ is a $(1, 2, 1)$ -scheme (in Section 4 you will see another equivalent condition for a scheme to be reduced). Any r -balanced scheme is obtained by the restriction of the tensor product of a reduced r -balanced scheme and a 1-balanced scheme (see Theorem 3.7). Now we focus our attention on reduced balanced schemes. It seems a quite difficult problem to find possible n such that there exists a reduced (m, n, r) -scheme for given m and r . Actually, D. G. Higman asked if there exists a reduced $(m, 3, 3)$ -scheme for some m (see [9, Section 8, p.229]). Furthermore, H. Weilandt conjectured that a permutation group of prime degree has at most two inequivalent permutation representations (see [1]). The following theorem shows that $n = 1$ is a unique case under certain assumptions.

Theorem 1.3. *Let $\mathcal{C}$ be a reduced (m, n, r) -scheme and p a prime. Then we have the following:*

- (i) If $m < 2r$, then $n = 1$.
- (ii) If $p \nmid m$ and $\mathcal{C}_X$ is p -valanced for some $X \in \text{Fib}(\mathcal{C})$, then $n = 1$.

The preceding theorem is applied to characterize (m, n, r) -schemes up to $m \leq 11$ as follows.

Theorem 1.4. *If $\mathcal{C}$ is a reduced (m, n, r) -scheme and $m \leq 11$, then $n \leq 2$.*

Let us show the organization of this article. In Section 2 we prepare some terminologies related to schemes. Section 3 is devoted to balanced schemes. First we investigate the features of balanced schemes. Indeed, we shall characterize a balanced scheme in terms of its central primitive idempotents and we prove Theorem 1.1. Secondly we shall characterize schemes with at most two central primitive idempotents and we prove Theorem 1.2. In Section 4 we shall extend the notion of inequivalent permutation representations to schemes. Namely, we shall define reduced (m, n, r) -schemes and then introduce some examples and known constructions of them to support our theory. Finally in Section 5, first we prove Theorem 1.3, secondly we shall enumerate reduced (m, n, r) -schemes for $m \leq 11$ in order to prove Theorem 1.4.

2. Preliminaries

According to [3] we prepare some terminologies related to schemes. For the remainder of this section we assume that $\mathcal{C} = (V, \mathcal{R})$ is a scheme. One can see that $V = \bigcup_{X \in \text{Fib}(\mathcal{C})} X$ (disjoint union) and

$$\mathcal{R} = \bigcup_{X, Y \in \text{Fib}(\mathcal{C})} \mathcal{R}_{X, Y} \quad (\text{disjoint union}), \quad (1)$$

where $\mathcal{R}_{X, Y} := \{R \in \mathcal{R} \mid R \subseteq X \times Y\}$. We shall denote $\mathcal{R}_{X, X}$ by $\mathcal{R}_X$.

Let $X, Y \in \text{Fib}(\mathcal{C})$ and R be a non-empty union of basis relations in $\mathcal{R}_{X, Y}$. For $(x, y) \in R$ we set $R_{\text{out}}(x) = \{u \mid (x, u) \in R\}$ and $R_{\text{in}}(y) = \{v \mid (v, y) \in R\}$. We shall denote the size of $R_{\text{out}}(x)$ and that of $R_{\text{in}}(y)$ by d_R and e_R , respectively. It is easy to see that

$$|X|d_R = |R| = |Y|e_R. \quad (2)$$

For $\mathcal{D} \subseteq \mathcal{R}$ we define $d_{\mathcal{D}} := \sum_{R \in \mathcal{D}} d_R$ as well as $e_{\mathcal{D}} := \sum_{R \in \mathcal{D}} e_R$. For instance $d_{\mathcal{R}_{X, Y}} = |Y|$ and $e_{\mathcal{R}_{X, Y}} = |X|$. For all $X, Y \in \text{Fib}(\mathcal{C})$, we define the multi-set $d_{X, Y} := \{d_R \mid R \in \mathcal{R}_{X, Y}\}$.

Note that $d_R = e_R$ for each $R \in \mathcal{R}$ if and only if $|X| = |Y|$ for all $X, Y \in \text{Fib}(\mathcal{C})$. A scheme $\mathcal{C}$ is called *half-homogeneous* if the latter condition holds. If $\mathcal{C}$ is a half-homogeneous scheme, then $d_R (= e_R)$ is called the *degree* or the *valency* of R . Given a prime p a half-homogeneous scheme $\mathcal{C}$ is called *p-valenced* if the degree of each basis relation of $\mathcal{C}$ is a power of p .

A basis relation $R \in \mathcal{R}$ is called *thin* if $d_R = e_R = 1$ and a scheme $\mathcal{C}$ is called a *homogeneous scheme* or (*association scheme*) if $|\text{Fib}(\mathcal{C})| = 1$ or

equivalently, if $\Delta_V \in \mathcal{R}$ (for more details regarding association schemes we refer to [15]). Given $X \in \text{Fib}(\mathcal{C})$ the pair $\mathcal{C}_X = (X, \mathcal{R}_X)$ is a homogeneous scheme called the *homogeneous component* of $\mathcal{C}$ corresponding to X .

Definition 2.1. For each $R \in \mathcal{R}$ we define a $\{0, 1\}$ -matrix A_R whose rows and columns are simultaneously indexed by the elements of V such that the (u, v) -entry of A_R is one if and only if $(u, v) \in R$. Then A_R is called the *adjacency matrix* of R . Note that the subspace of $\text{Mat}_V(\mathbb{C})$ spanned by $\{A_R \mid R \in \mathcal{R}\}$ is a subalgebra, called the *adjacency algebra* of $\mathcal{C}$ and denoted by $\mathcal{A}(\mathcal{C})$.

Let $\mathcal{A}$ be the adjacency algebra of $\mathcal{C} = (V, \mathcal{R})$. Then the set $\{A_R \mid R \in \mathcal{R}\}$ which is a basis of $\mathcal{A}$ satisfies the following conditions:

- (C'1) $J_V = \sum_{R \in \mathcal{R}} A_R \in \mathcal{A}$ where J_V is the matrix whose entries are all one.
- (C'2) The identity matrix $I_V \in \mathcal{A}$.
- (C'3) $A_{R^t} = A_R^t$ for every $R \in \mathcal{R}$ where A_R^t is the transpose of A_R .
- (C'4) For every $R, S \in \mathcal{R}$, $A_R A_S = \sum_{T \in \mathcal{R}} c_{RS}^T A_T$.

A scheme is called *trivial* if all its fibers are singletons. We denote a trivial scheme on n points by $\mathcal{T}_n$. Note that $\mathcal{A}(\mathcal{T}_n) \cong \text{Mat}_n(\mathbb{C})$ and it is easy to see that a scheme is trivial if and only if it is 1-balanced.

By $\text{Fib}^*(\mathcal{C})$ we mean the set of all non-empty unions of fibres of $\mathcal{C}$. Given $U \in \text{Fib}^*(\mathcal{C})$ we set $\mathcal{R}_U := \{R_U \mid R \in \mathcal{R}\}$ where $R_U = R \cap (U \times U)$. Then the pair $\mathcal{C}_U = (U, \mathcal{R}_U)$ is a scheme on U called the *restriction* of $\mathcal{C}$ to U . Note that $\mathcal{C}_U$ is homogeneous whenever $U \in \text{Fib}(\mathcal{C})$.

Given $U, U' \in \text{Fib}^*(\mathcal{C})$ we define $\mathcal{A}_{U, U'}$ to be the subspace of $\mathcal{A}$ spanned by the set $\{A_R \mid R \in \mathcal{R}, R \subseteq U \times U'\}$. Then the following hold:

- (i) Given $U \in \text{Fib}^*(\mathcal{C})$ we have $\mathcal{A}_U = I_U \mathcal{A} I_U \cong \mathcal{A}(\mathcal{C}_U)$ where $I_U := \sum_{\substack{X \in \text{Fib}(\mathcal{C}), \\ X \subseteq U}} A_{\Delta_X}$.
- (ii) For all $X, Y, Z, W \in \text{Fib}(\mathcal{C})$, $\mathcal{A}_{X, Y} \mathcal{A}_{Z, W} \subseteq \delta_{YZ} \mathcal{A}_{X, W}$.
- (iii) $\mathcal{A} = \bigoplus_{X, Y \in \text{Fib}(\mathcal{C})} \mathcal{A}_{X, Y}$.

A basis relation S of $\mathcal{C}$ is called *symmetric* if $S^t = S$ and $\mathcal{C}$ is called *symmetric* if each basis relation of $\mathcal{C}$ is symmetric; and $\mathcal{C}$ is called *commutative* if $c_{RS}^T = c_{SR}^T$ for all $R, S, T \in \mathcal{R}$. This is equivalent to $A_R A_S = A_S A_R$ for all

$R, S \in \mathcal{R}$. It is known that symmetric schemes are commutative and that the converse does not hold. Furthermore, one can see that a commutative scheme is a homogeneous one.

Lemma 2.1 ([8, (4.2)], [15, Theorem 4.5.1]). *If $\mathcal{C} = (V, \mathcal{R})$ is a homogeneous scheme and $|\mathcal{R}| \leq 5$, then $\mathcal{C}$ is commutative.*

Given $R, S \in \mathcal{R}$ the *complex product* of them is defined to be $RS = \{T \in \mathcal{R} \mid c_{RS}^T > 0\}$ and the *relational product* $R \circ S$ is defined as follows.

$$R \circ S := \{(u, v) \mid \exists w \in V; (u, w) \in R, (w, v) \in S\}.$$

Note that $R \circ S = \bigcup_{T \in RS} T$ and $d_{R \circ S} = d_{RS}$.

Lemma 2.2. *Let $\mathcal{C}$ be a scheme and $X, Y, Z \in \text{Fib}(\mathcal{C})$. Then for all $R \in \mathcal{R}_{X,Y}$, $S \in \mathcal{R}_{Y,Z}$ and $T \in \mathcal{R}_{X,Z}$ the following hold:*

- (i) $d_R d_S = \sum_{T \in \mathcal{R}_{X,Z}} c_{RS}^T d_T$.
- (ii) $c_{RS}^T d_T = c_{TS^t}^R d_R = c_{R^t T}^S d_S$ and $\text{lcm}(d_R, d_S) \mid c_{RS}^T d_T$.
- (iii) $d_R = \sum_{S \in \mathcal{R}_{Y,Z}} c_{RS}^T$, $e_R = \sum_{S \in \mathcal{R}_{Y,Z}} c_{R^t T}^S$, $c_{RS}^T \leq \min\{d_R, e_S\}$ and $R\mathcal{R}_{Y,Z} = \mathcal{R}_{X,Z}$.
- (iv) $d_R \delta_{SR^t} = c_{RS}^{\Delta_X}$ and $e_R \delta_{SR^t} = c_{SR}^{\Delta_Y}$ where δ denotes the Kronecker's delta.
- (v) $d_S \leq d_{RS} \leq d_R d_S$ and $e_R \leq e_{RS} \leq e_R e_S$.
- (vi) If $RS = T$ and $d_T \leq d_R$, then $R = TS^t$.
- (vii) If $d_R = 2$, then $RR^t = \{\Delta_X, S\}$ where S is a symmetric basis relation of $\mathcal{R}_X$ with $d_S \leq 2$.
- (viii) $|RS| \leq \gcd(d_R, d_S)$.

Proof. The proof is done by the same procedure as [15, Lemma 1.4.2, 1.4.3, 1.5.2, 1.5.3, 1.5.6]. $\square$

Lemma 2.3. *Let $S \in \mathcal{R}_{X,Y}$ and $L_S := \{R \in \mathcal{R}_X \mid RS = \{S\}\}$. Then*

$$d_{L_S} \mid \gcd(|X|, e_S).$$

Proof. Let $y \in Y$ and $x \in S_{in}(y)$. The condition $RS = \{S\}$ shows that $\bigcup_{R \in L_S} R_{in}(x) \subseteq S_{in}(y)$ and $\bigcup_{R \in L_S} R$ is an equivalence relation on X . Since $y \in Y$ and $x \in S_{in}(y)$ are arbitrarily taken, all equivalence classes have the same size d_{L_S} . It follows that d_{L_S} divides both d_S and $|X|$. $\square$

Lemma 2.4. *Let $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$ and $R, S \in \mathcal{R}_{X,Y}$ with $R \neq S$. Then $T \in R^t R \cap S^t S$ for some $T \in \mathcal{R}_Y$ with $T \neq \Delta_Y$ if and only if $c_{RS^t}^{T'} \geq 2$ for some $T' \in \mathcal{R}_X$.*

Proof. Let us prove the necessity. By the assumption $c_{R^t R}^T \neq 0$ and $c_{S^t S}^T \neq 0$. Taking $(y, y') \in T$ (of course $y \neq y'$) there exist $x, x' \in X$ such that $(x, y), (x, y') \in R$ and $(x', y), (x', y') \in S$. On the other hand, there exists $T' \in \mathcal{R}_X$ such that $(x, x') \in T'$. It follows that $c_{RS^t}^{T'} \geq 2$ (see the Figure 1). Sufficiency follows from Figure 1 since $c_{RS^t}^{T'} \geq 2$ implies that $y \neq y'$.




Figure 1.

□

Let $U, U' \in \text{Fib}^*(\mathcal{C})$ such that $U \cap U' = \emptyset$ and $V = U \cup U'$. Then we say that $\mathcal{C}$ is the *internal direct sum* of $\mathcal{C}_U$ and $\mathcal{C}_{U'}$ if $|\mathcal{R}_{X,Y}| = 1$ for all $X, Y \in \text{Fib}(\mathcal{C})$ with $X \subseteq U$ and $Y \subseteq U'$. In this case we shall write $\mathcal{C} = \mathcal{C}_U \boxplus \mathcal{C}_{U'}$.

Let $\mathcal{C}_i = (V_i, \mathcal{R}_i)$ ($i = 1, 2$) be schemes.

We set

$$\mathcal{R}_1 \otimes \mathcal{R}_2 = \{R_1 \otimes R_2 \mid R_1 \in \mathcal{R}_1, R_2 \in \mathcal{R}_2\},$$

where $R_1 \otimes R_2 = \{((u_1, u_2), (v_1, v_2)) \mid (u_1, v_1) \in R_1, (u_2, v_2) \in R_2\}$. Then $\mathcal{C} = (V_1 \times V_2, \mathcal{R}_1 \otimes \mathcal{R}_2)$ is a scheme called the *tensor product* of $\mathcal{C}_1$ and $\mathcal{C}_2$ and denoted by $\mathcal{C}_1 \otimes \mathcal{C}_2$. One can see that $\text{Fib}(\mathcal{C}) = \text{Fib}(\mathcal{C}_1) \times \text{Fib}(\mathcal{C}_2)$.

An *isomorphism* from $\mathcal{C}_1$ to $\mathcal{C}_2$ is defined to be a bijection $\psi : V_1 \cup \mathcal{R}_1 \rightarrow V_2 \cup \mathcal{R}_2$ such that for all $u, v \in V_1$ and $R \in \mathcal{R}_1$, $(u, v) \in R$ if and only if $(\psi(u), \psi(v)) \in \psi(R)$. We say that $\mathcal{C}_1$ is *isomorphic* to $\mathcal{C}_2$ and denote it by $\mathcal{C}_1 \simeq \mathcal{C}_2$ if there exists an isomorphism from $\mathcal{C}_1$ to $\mathcal{C}_2$.

Let $\mathcal{A}$ be the adjacency algebra of $\mathcal{C}$. Since $\mathcal{A}$ is closed under the complex conjugate transpose map, $\mathcal{A}$ is semisimple. By the Wedderburn theorem $\mathcal{A}$ is isomorphic to a direct sum of full matrix algebras over $\mathbb{C}$:

$$\mathcal{A} = \bigoplus_{P \in \mathcal{P}(\mathcal{C})} \mathcal{A}P \cong \bigoplus_{P \in \mathcal{P}(\mathcal{C})} \text{Mat}_{n_P}(\mathbb{C}), \quad (3)$$

where $\mathcal{P}(\mathcal{C})$ is the set of central primitive idempotents of $\mathcal{A}$, n_P is a positive integer and $\text{Mat}_{n_P}(\mathbb{C})$ is the full matrix algebra of complex $n_P \times n_P$ matrices. A comparison of dimensions of the left- and right-hand sides of (3) shows that

$$|\mathcal{R}| = \sum_{P \in \mathcal{P}(\mathcal{C})} n_P^2. \quad (4)$$

It is known that $\mathcal{C}$ is commutative if and only if $n_P = 1$ for each $P \in \mathcal{P}(\mathcal{C})$. Denote by $\mathbb{C}^V$ the natural $\mathcal{A}$ -module spanned by the elements of V . As $I_V = \sum_{P \in \mathcal{P}(\mathcal{C})} P$ we have

$$\mathbb{C}^V = \bigoplus_{P \in \mathcal{P}(\mathcal{C})} P\mathbb{C}^V. \quad (5)$$

For each $P \in \mathcal{P}(\mathcal{C})$ we set $m_P := \dim_{\mathbb{C}}(P\mathbb{C}^V)/n_P$. Then the decomposition (5) shows that

$$|V| = \sum_{P \in \mathcal{P}(\mathcal{C})} m_P n_P. \quad (6)$$

The numbers m_P and n_P are called the *multiplicity* and the *degree* of P . Set $P_0 = \sum_X J_X/|X|$ where X runs over the fibers of the scheme $\mathcal{C}$ and $J_X = \sum_{R \in \mathcal{R}_X} A_R$. Note that P_0 is a central primitive idempotent of the algebra $\mathcal{A}$, which is called *principal*. It is known that

$$(m_{P_0}, n_{P_0}) = (1, |\text{Fib}(\mathcal{C})|). \quad (7)$$

If $\mathcal{C}$ is homogeneous, then $P_0 = J_V/|V|$ and $m_{P_0} = n_{P_0} = 1$.

Below for $X \in \text{Fib}^*(\mathcal{C})$ and $P \in \mathcal{P}(\mathcal{C})$ put $P_X = PI_X$ and set

$$\mathcal{P}_X(\mathcal{C}) = \left\{ P \in \mathcal{P}(\mathcal{C}) \mid P_X \neq 0 \right\} \quad \text{and} \quad \text{Supp}(P) = \left\{ X \in \text{Fib}(\mathcal{C}) \mid P_X \neq 0 \right\}.$$

Theorem 2.5 ([4, Proposition 2.1]). *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. Then the following hold:*

- (i) *For each $X \in \text{Fib}^*(\mathcal{C})$ the mapping $P \mapsto P_X$ induces a bijection between $\mathcal{P}_X(\mathcal{C})$ and $\mathcal{P}(\mathcal{C}_X)$.*

- (ii) For all $P \in \mathcal{P}(\mathcal{C})$ and $X \in \text{Supp}(P)$, $n_P = \sum_{X \in \text{Supp}(P)} n_{P_X}$ and $m_P = m_{P_X}$.

Lemma 2.6. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. Then the following hold:*

- (i) $\mathcal{P}(\mathcal{C}) = \mathcal{P}_X(\mathcal{C})$ for each $X \in \text{Fib}(\mathcal{C})$ if and only if $\text{Supp}(P) = \text{Fib}(\mathcal{C})$ for each $P \in \mathcal{P}(\mathcal{C})$.
(ii) $\text{Supp}(P) \neq \emptyset$ for each $P \in \mathcal{P}(\mathcal{C})$, and

$$\mathcal{P}(\mathcal{C}) = \bigcup_{X \in \text{Fib}(\mathcal{C})} \mathcal{P}_X(\mathcal{C}). \quad (8)$$

Besides, $\mathcal{P}(\mathcal{C}) = \mathcal{P}_U(\mathcal{C}) \cup \mathcal{P}_{U'}(\mathcal{C})$ where $U, U' \in \text{Fib}^*(\mathcal{C})$ with $U \cap U' = \emptyset$ and $V = U \cup U'$.

Proof. (i) Let $X \in \text{Fib}(\mathcal{C})$ and $P \in \mathcal{P}(\mathcal{C})$. Then $P \in \mathcal{P}_X(\mathcal{C})$ if and only if $X \in \text{Supp}(P)$. This completes the proof.

(ii) Let $P \in \mathcal{P}(\mathcal{C})$ such that $\text{Supp}(P) = \emptyset$. Then for all $X \in \text{Fib}(\mathcal{C})$, $PI_X = 0$ and then $P = PI_V = \sum_{X \in \text{Fib}(\mathcal{C})} PI_X = 0$, a contradiction. Therefore, $\text{Supp}(P) \neq \emptyset$. Let $P \in \mathcal{P}(\mathcal{C})$, as $\text{Supp}(P) \neq \emptyset$, there exists $X \in \text{Fib}(\mathcal{C})$ such that $PI_X \neq 0$. This means that $P \in \mathcal{P}_X(\mathcal{C})$ and the proof of (8) is completed.

Let $P \in \mathcal{P}(\mathcal{C})$. Then $P \in \mathcal{P}_X(\mathcal{C})$ for some $X \in \text{Fib}(\mathcal{C})$. Since $V = U \cup U'$, $X \subseteq U$ or $X \subseteq U'$. It follows that $P \in \mathcal{P}_U(\mathcal{C})$ or $P \in \mathcal{P}_{U'}(\mathcal{C})$. This completes the proof. $\square$

Proposition 2.7 ([9, p.223], [8, p.22 (8.1)]). *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. Then the following hold:*

- (i) Let $X, Y \in \text{Fib}^*(\mathcal{C})$ such that $X \cap Y = \emptyset$ and $V = X \cup Y$. Then

$$\dim_{\mathbb{C}}(\mathcal{A}_{X,Y}) = \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y}.$$

- (ii) For all $X, Y \in \text{Fib}(\mathcal{C})$, $|\mathcal{R}_{X,Y}| = \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y}.$

Lemma 2.8. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme with the adjacency algebra $\mathcal{A}(\mathcal{C})$. If $U, U' \in \text{Fib}^*(\mathcal{C})$ such that $U \cap U' = \emptyset$, then*

$$|\text{Fib}(\mathcal{C}_U)| |\text{Fib}(\mathcal{C}_{U'})| \leq \dim_{\mathbb{C}}(\mathcal{A}_{U,U'}). \quad (9)$$

Furthermore, the equality holds if and only if $\mathcal{C}_{U \cup U'} = \mathcal{C}_U \boxplus \mathcal{C}_{U'}$.

Proof. It is clear that $\text{Fib}(\mathcal{C}_U) = \{X \in \text{Fib}(\mathcal{C}) \mid X \subseteq U\}$ as well as $\text{Fib}(\mathcal{C}_{U'}) = \{Y \in \text{Fib}(\mathcal{C}) \mid Y \subseteq U'\}$. By the assumption,

$$\mathcal{A}_{U,U'} = \bigoplus_{\substack{X \in \text{Fib}(\mathcal{C}_U), \\ Y \in \text{Fib}(\mathcal{C}_{U'})}} \mathcal{A}_{X,Y}.$$

On the other hand, $\dim_{\mathbb{C}}(\mathcal{A}_{X,Y}) = |\mathcal{R}_{X,Y}| \geq 1$ for each $X \in \text{Fib}(\mathcal{C}_U)$ and $Y \in \text{Fib}(\mathcal{C}_{U'})$. It follows that

$$|\text{Fib}(\mathcal{C}_U)| |\text{Fib}(\mathcal{C}_{U'})| \leq \sum_{\substack{X \in \text{Fib}(\mathcal{C}_U), \\ Y \in \text{Fib}(\mathcal{C}_{U'})}} |\mathcal{R}_{X,Y}| = \dim_{\mathbb{C}}(\mathcal{A}_{U,U'}).$$

The equality holds if and only if $|\mathcal{R}_{X,Y}| = 1$ for all $X \in \text{Fib}(\mathcal{C}_U)$ and $Y \in \text{Fib}(\mathcal{C}_{U'})$ which is exactly the definition of internal direct sums. $\square$

Lemma 2.9. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme with the principal idempotent P_0 and let $U, U' \in \text{Fib}^*(\mathcal{C})$ such that $U \cap U' = \emptyset$ and $V = U \cup U'$. Then $\mathcal{C} = \mathcal{C}_U \boxplus \mathcal{C}_{U'}$ if and only if $\mathcal{P}_U(\mathcal{C}) \cap \mathcal{P}_{U'}(\mathcal{C}) = \{P_0\}$.*

Proof. Let us prove the sufficiency first. It is clear that $P_0 \in \mathcal{P}_U(\mathcal{C}) \cap \mathcal{P}_{U'}(\mathcal{C})$. By Lemma 2.8 and Proposition 2.7 (i) we have

$$|\text{Fib}(\mathcal{C}_U)| |\text{Fib}(\mathcal{C}_{U'})| = \sum_{P \in \mathcal{P}_U \cap \mathcal{P}_{U'}} n_P n_{P_{U'}}.$$

Since $n_{P_0U} = |\text{Fib}(\mathcal{C}_U)|$ and $n_{P_0U'} = |\text{Fib}(\mathcal{C}_{U'})|$, it follows that

$$\mathcal{P}_U(\mathcal{C}) \cap \mathcal{P}_{U'}(\mathcal{C}) = \{P_0\}. \quad (10)$$

Conversely, if $\mathcal{P}_U(\mathcal{C}) \cap \mathcal{P}_{U'}(\mathcal{C}) = \{P_0\}$, then by Proposition 2.7 (i),

$$\dim_{\mathbb{C}}(\mathcal{A}_{U,U'}) = |\text{Fib}(\mathcal{C}_U)| |\text{Fib}(\mathcal{C}_{U'})|.$$

It follows from Lemma 2.8 that $\mathcal{C} = \mathcal{C}_U \boxplus \mathcal{C}_{U'}$. $\square$

3. Characterization of balanced schemes

Proof of Theorem 1.1:

First we prove the necessity. Let $X, Y \in \text{Fib}(\mathcal{C})$. By Proposition 2.7, $|\mathcal{R}_{X,Y}| = \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y}$. By the Cauchy-Schwarz inequality we have

$$\begin{aligned} |\mathcal{R}_{X,Y}|^2 &= \left(\sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y} \right)^2 \leq \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X}^2 \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_Y}^2 \\ &\leq \sum_{P \in \mathcal{P}_X} n_{P_X}^2 \sum_{P \in \mathcal{P}_Y} n_{P_Y}^2 \\ &= |\mathcal{R}_X| |\mathcal{R}_Y| = |\mathcal{R}_{X,Y}|^2. \end{aligned}$$

This implies that

$$\left(\sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y} \right)^2 = \sum_{P \in \mathcal{P}_X} n_{P_X}^2 \sum_{P \in \mathcal{P}_Y} n_{P_Y}^2.$$

It follows that $\mathcal{P}_X(\mathcal{C}) = \mathcal{P}_Y(\mathcal{C})$ and thus applying Lemma 2.6 (i) we have $\mathcal{P}(\mathcal{C}) = \mathcal{P}_X(\mathcal{C})$. Consequently, the mapping $\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C}_X)$ ($P \mapsto P_X$) is well-defined and bijective by Theorem 2.5. Since the equality holds in the Cauchy-Schwarz inequality, we have $\langle n_{P_X} | P \in \mathcal{P}(\mathcal{C}) \rangle = \alpha \langle n_{P_Y} | P \in \mathcal{P}(\mathcal{C}) \rangle$. However, $\alpha = 1$ since $|\mathcal{R}_X| = |\mathcal{R}_Y|$. Hence, $n_{P_X} = n_{P_Y}$ for all $P \in \mathcal{P}(\mathcal{C})$. Therefore, by Theorem 2.5 and Lemma 2.6 (ii),

$$n_P = \sum_{X \in \text{Supp}(P)} n_{P_X} = \sum_{X \in \text{Fib}(\mathcal{C})} n_{P_X} = |\text{Fib}(\mathcal{C})| n_{P_X}.$$

Now let us prove the sufficiency. Given $X, Y \in \text{Fib}(\mathcal{C})$ the assumption along with Theorem 2.5 assert that $\mathcal{P}(\mathcal{C}) = \mathcal{P}_X(\mathcal{C}) = \mathcal{P}_Y(\mathcal{C})$ and $n_{P_X} = n_{P_Y}$ for each $P \in \mathcal{P}(\mathcal{C})$. On the other hand, by Proposition 2.7 (ii), we have

$$|\mathcal{R}_{X,Y}| = \sum_{P \in \mathcal{P}_X \cap \mathcal{P}_Y} n_{P_X} n_{P_Y} = \sum_{P \in \mathcal{P}(\mathcal{C})} n_{P_X}^2 = |\mathcal{R}_X|.$$

Hence, $\mathcal{C}$ is balanced. □

Corollary 3.1. *Let $\mathcal{C}$ be an r -balanced scheme. If $\mathcal{C}_X$ is commutative for some $X \in \text{Fib}(\mathcal{C})$, then $|\mathcal{P}(\mathcal{C})| = r$. For instance, $|\mathcal{P}(\mathcal{C})| = r$ if $r \leq 5$.*

Proof. Let $X \in \text{Fib}(\mathcal{C})$. Since $\mathcal{C}_X$ is commutative, $|\mathcal{P}(\mathcal{C}_X)| = |\mathcal{R}_X| = r$. By Theorem 1.1, $|\mathcal{P}(\mathcal{C})| = |\mathcal{P}(\mathcal{C}_X)| = r$. In particular, if $r \leq 5$, then by Lemma 2.1, $\mathcal{C}_X$ is commutative and thus $|\mathcal{P}(\mathcal{C})| = r$. $\square$

Proof of Theorem 1.2 (i):

Let $X \in \text{Fib}(\mathcal{C})$. By Theorem 2.5, $|\mathcal{P}(\mathcal{C}_X)| = 1$. On the other hand, $\mathcal{C}_X = (X, \mathcal{R}_X)$ is a homogeneous scheme, so $|X| = m_{P_0X} n_{P_0X} = 1$, by (7). Hence, every fiber of $\mathcal{C}$ is a singleton and thus $\mathcal{C}$ is trivial. Conversely, the adjacency algebra of a trivial scheme is the full matrix algebra and thus it has only one central primitive idempotent. $\square$

In order to prove Theorem 1.2 (ii), we need the following preparations.

Lemma 3.2 ([8, (4.2)]). *Let $\mathcal{C} = (V, \mathcal{R})$ be a homogenous scheme. Then $|\mathcal{P}(\mathcal{C})| = 2$ if and only if $|\mathcal{R}| = 2$.*

Lemma 3.3. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. If $\mathcal{P}(\mathcal{C}) = \{P_0, P_1\}$ with $P_0 \neq P_1$, then the following hold:*

- (i) $X \notin \text{Supp}(P_1)$ if and only if $|X| = 1$.
- (ii) $|\mathcal{R}_X| = \begin{cases} 2 & \text{if } X \in \text{Supp}(P_1) \\ 1 & \text{if } X \notin \text{Supp}(P_1) \end{cases}$.

Proof. (i) Since $I_V = P_0 + P_1$, $P_1 = \sum_{X \in \text{Fib}(\mathcal{C})} (I_X - J_X/|X|)$. Let $X \in \text{Fib}(\mathcal{C})$. Then $X \notin \text{Supp}(P_1)$ if and only if $0 = P_1 I_X = I_X - J_X/|X|$ if and only if $|X| = 1$.

(ii) If $X \in \text{Supp}(P_1)$, then $P_1 I_X \neq 0$ and by Theorem 2.5, $|\mathcal{P}(\mathcal{C}_X)| = 2$. Since $\mathcal{C}_X = (X, \mathcal{R}_X)$ is homogeneous, it follows from Lemma 3.2 that $|\mathcal{R}_X| = 2$. If $X \notin \text{Supp}(P_1)$, then by (i), we have $|X| = 1$. It follows that $|\mathcal{R}_X| = 1$. $\square$

Lemma 3.4. *Let $\mathcal{C} = (V, \mathcal{R})$ be a scheme. If $\mathcal{P}(\mathcal{C}) = \{P_0, P_1\}$ with $P_0 \neq P_1$, then the following hold:*

- (i) $n_{P_1} = |\text{Supp}(P_1)|$ and $|X| = 1 + m_{P_1}$ for each $X \in \text{Supp}(P_1)$.
- (ii) $|\mathcal{R}_{X,Y}| = 2$ for each $X, Y \in \text{Supp}(P_1)$.

Proof. (i) Let $X \in \text{Supp}(P_1)$. By Lemma 3.3, $|\mathcal{R}_X| = 2$ and thus by Lemma 2.1, $\mathcal{C}_X$ is commutative. By Theorem 2.5 we have

$$n_{P_1} = \sum_{X \in \text{Supp}(P_1)} n_{P_1X} = |\text{Supp}(P_1)|.$$

Thus (6) implies that $|X| = 1 + m_{P_1}$.

(ii) Let $X, Y \in \text{Supp}(P_1)$. Then by Lemma 3.3 (ii), $|\mathcal{R}_Y| = |\mathcal{R}_X| = 2$. It follows from Lemma 3.2 that $\mathcal{P}_X(\mathcal{C}) \cap \mathcal{P}_Y(\mathcal{C}) = \mathcal{P}(\mathcal{C})$. Therefore, Proposition 2.7 (ii) implies that $|\mathcal{R}_{X,Y}| = 2$. $\square$

Proof of Theorem 1.2 (ii): Let $\mathcal{P}(\mathcal{C}) = \{P_0, P_1\}$ and set $U := \bigcup_{X \in \text{Supp}(P_1)} X$ and $U' := V \setminus U$. If $X \in \text{Supp}(P_1)$ and $Y \notin \text{Supp}(P_1)$, then $|\mathcal{R}_{X,Y}| = 1$, since $|Y| = 1$ by Lemma 3.3. This implies that

$$\mathcal{C} = \mathcal{C}_U \boxplus \mathcal{C}_{U'}.$$

Note that by Lemma 3.3 (ii) and Lemma 3.4 (ii), $\mathcal{C}_U$ is 2-balanced and $\mathcal{C}_{U'}$ is 1-balanced. Conversely, by Lemma 2.9 and Corollary 3.1, $|\mathcal{P}(\mathcal{C})| = |\mathcal{P}(\mathcal{C}_1 \boxplus \mathcal{C}_2)| = |\mathcal{P}(\mathcal{C}_1)| + |\mathcal{P}(\mathcal{C}_2)| - 1 = |\mathcal{P}(\mathcal{C}_2)| = 2$. $\square$

Corollary 3.5. *If $\mathcal{C}$ is a balanced scheme, then the following hold:*

- (i) $\mathcal{C}$ is half-homogeneous.
- (ii) For every $X, Y \in \text{Fib}(\mathcal{C})$, $\mathcal{A}_X$ and $\mathcal{A}_Y$ are isomorphic as $\mathbb{C}$ -algebras.

Proof. (i) Let $X \in \text{Fib}(\mathcal{C})$ and consider the scheme $\mathcal{C}_X = (X, \mathcal{R}_X)$. It follows from Theorem 1.1 that the mapping $\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C}_X)$ ($P \mapsto P_X$) is bijective with $n_P = |\text{Fib}(\mathcal{C})| n_{P_X}$. By (6) and Theorem 2.5 (ii), the size of X is computed as follows.

$$|X| = \sum_{P \in \mathcal{P}(\mathcal{C})} n_{P_X} m_{P_X} = \frac{1}{|\text{Fib}(\mathcal{C})|} \sum_{P \in \mathcal{P}(\mathcal{C})} n_P m_P = \frac{|V|}{|\text{Fib}(\mathcal{C})|}.$$

Hence, the size of each fiber is constant and thus $\mathcal{C}$ is half-homogeneous.

(ii) Let $X, Y \in \text{Fib}(\mathcal{C})$ and $P \in \mathcal{P}(\mathcal{C})$. Then by Theorem 1.1, $n_{P_X} = n_{P_Y}$. It follows from (3) that

$$\mathcal{A}_X = \bigoplus_{P \in \mathcal{P}(\mathcal{C})} \text{Mat}_{n_{P_X}}(\mathbb{C}) \cong \bigoplus_{P \in \mathcal{P}(\mathcal{C})} \text{Mat}_{n_{P_Y}}(\mathbb{C}) = \mathcal{A}_Y.$$

$\square$

Given a scheme $\mathcal{C}$ we define a relation $E_{\mathcal{C}}$ on $\text{Fib}(\mathcal{C})$ as follows.

$$E_{\mathcal{C}} := \left\{ (X, Y) \in \text{Fib}(\mathcal{C}) \mid \exists R \in \mathcal{R}_{X,Y} ; d_R = e_R = 1 \right\}. \quad (11)$$

Lemma 3.6. $E_{\mathcal{C}}$ is an equivalence relation on $\text{Fib}(\mathcal{C})$.

Proof. For each $X \in \text{Fib}(\mathcal{C})$, Δ_X is a thin basis relation in $\mathcal{R}_X$ and thus $E_{\mathcal{C}}$ is reflexive. If $R \in \mathcal{R}_{X,Y}$ is thin, then $R^t \in \mathcal{R}_{Y,X}$ is also thin and then $E_{\mathcal{C}}$ is symmetric. Let $X, Y, Z \in \text{Fib}(\mathcal{C})$ and $R \in \mathcal{R}_{X,Y}, S \in \mathcal{R}_{Y,Z}$ such that $d_R = d_S = 1$ and $e_R = e_S = 1$. It follows from Lemma 2.2 (v) that RS is a thin basis relation in $\mathcal{R}_{X,Z}$ and thus $E_{\mathcal{C}}$ is transitive. $\square$

Theorem 3.7. Let $\mathcal{C}$ be an (m, n, r) -scheme with the equivalence relation $E_{\mathcal{C}}$. If $\mathfrak{X}$ is a transversal of $E_{\mathcal{C}}$ in $\text{Fib}(\mathcal{C})$, then $\mathcal{C}$ is a restriction of $\mathcal{C}_{U_{\mathfrak{X}}} \otimes \mathcal{T}_n$ where $U_{\mathfrak{X}} = \bigcup_{X \in \mathfrak{X}} X$ and $\mathcal{T}_n$ is a $(1, n, 1)$ -scheme.

Proof. Let $I_n := \{1, \dots, n\}$ and $E_n := \{e_{ij} \mid 1 \leq i, j \leq n\}$ where for $1 \leq i, j \leq n$, $e_{ij} = \{(i, j)\}$. Then $\mathcal{T}_n = (I_n, E_n)$. Let $\mathfrak{X} = \{X_1, \dots, X_s\}$ and suppose that for each $i \in \{1, 2, \dots, s\}$, $E_{\mathcal{C}}(X_i) = \{X_{i1}, X_{i2}, \dots, X_{im_i}\}$ where $X_{i1} := X_i$ and X_{ij} 's are distinct fibers. In this case, $V = \bigcup_{i=1}^s \bigcup_{j=1}^{m_i} X_{ij}$. For all $i \in \{1, \dots, s\}$ and $j \in \{1, \dots, m_i\}$, there exists $R_{ij} \in \mathcal{R}_{X_{i1}, X_{ij}}$ with $d_{R_{ij}} = 1$. Therefore, there exists a bijection $R_{ij} : X_i \rightarrow X_{ij}, (x_i \mapsto x)$ where x is the unique element of X_{ij} such that $(x_i, x) \in R_{ij}$. Indeed, $R_{ij}(X_i) = X_{ij}$. Thus, for each $x \in V$, there exist unique $i \in \{1, \dots, s\}$ and $j \in \{1, \dots, m_i\}$ such that $R_{ij}(X_i) = X_{ij}$ and $x \in R_{ij}(X_i)$. We claim that the map ψ defined as follows is a monomorphism.

$$\begin{aligned} \psi : V \cup \mathcal{R} &\longrightarrow (U_{\mathfrak{X}} \times I_n) \cup (\mathcal{R}_{U_{\mathfrak{X}}} \otimes E_n). \\ x &\mapsto (x_i, j); \quad R_{ij}(x_i) = x, \\ R &\mapsto R_{ij} R R_{kl}^t \otimes e_{jl}; \quad R \in \mathcal{R}_{X_{ij}, X_{kl}} \end{aligned}$$

Note that ψ is injective, since R_{ij} is a bijection for all $i \in \{1, \dots, s\}$ and $j \in \{1, \dots, m_i\}$. Let $(x, y) \in R$ and $R \in \mathcal{R}_{X_{ij}, X_{kl}}$. Then there exists $(x_i, y_k) \in X_i \times X_k$ such that $R_{ij}(x_i) = x$ and $R_{kl}(y_k) = y$. This means that $(x_i, y_k) \in R_{ij} R R_{kl}^t$. It follows that, $(\psi(x), \psi(y)) = ((x_i, j), (y_k, l)) \in \psi(R)$. $\square$

The following is an immediate consequence of the preceding theorem.

Corollary 3.8. Let $\mathcal{C} = (V, \mathcal{R})$ be an (m, n, r) -scheme. Then $\mathcal{C} \simeq \mathcal{C}_X \otimes \mathcal{T}_n$ for $X \in \text{Fib}(\mathcal{C})$ if and only if $E_{\mathcal{C}}$ is trivial.

4. Reduced (m, n, r) -schemes

Definition 4.1. An (m, n, r) -scheme $\mathcal{C}$ is called *reduced* if its equivalence relation $E_{\mathcal{C}}$ is discrete.

Remark 4.1. Note that by Corollary 3.8, a balanced scheme $\mathcal{C}$ is reduced if and only if there exist no $X, Y \in \text{Fib}(\mathcal{C})$ such that $\mathcal{C}_{X \cup Y} \simeq \mathcal{C}_X \otimes \mathcal{T}_2$ where $\mathcal{T}_2$ is a $(1, 2, 1)$ -scheme.

Example 4.2. Let $(X, \mathcal{B}, \mathcal{I})$ be a symmetric design with the set X of points, the set $\mathcal{B}$ of blocks and the incidence relation $\mathcal{I} \subseteq X \times \mathcal{B}$. Set $V = X \cup \mathcal{B}$ (disjoint union) and define the relations R_i ($i = 1, \dots, 8$) on V as follows.

$$\begin{aligned} R_1 &= \Delta_X, & R_2 &= \Delta_{\mathcal{B}}, & R_3 &= (X \times X) \setminus \Delta_X, & R_4 &= (\mathcal{B} \times \mathcal{B}) \setminus \Delta_{\mathcal{B}}, \\ R_5 &= \mathcal{I}, & R_6 &= R_5^t, & R_7 &= (X \times \mathcal{B}) \setminus \mathcal{I}, & R_8 &= R_7^t. \end{aligned}$$

It is known that $(V, \{R_i\}_{i=1}^8)$ is an $(m, 2, 2)$ -scheme where $m = |X|$.

In [1] and [10] they mentioned $(m, n, 2)$ -schemes and $(m, 2, 3)$ -schemes as linked symmetric designs and strongly regular designs of the second kind, respectively.

Example 4.3 ([14], [1, p.6, Example (i)]). The split extension of the translation group of the vector space $\text{GF}(2^t)^{2k}$ by the symplectic group $\text{Sp}(2k, 2^t)$ has 2^t pairwise inequivalent doubly transitive representations of degree 2^{2kt} with the same character. This gives a reduced $(2^{2kt}, 2^t, 2)$ -scheme.

Lemma 4.2 ([2, Chapter 4]). Let $G = \text{PGL}(t, q)$ and Ω_k the set of k -dimensional subspaces of the vector space $\text{GF}(q)^t$. Let π_k denote the permutation character of G on Ω_k . Then we have the following:

For each $k \leq \frac{t}{2}$ there exist irreducible characters $\chi_0, \chi_1, \dots, \chi_k$ of G with $\chi_0 = 1_G$ such that

$$\pi_{t-k} = \pi_k = \sum_{i=0}^k \chi_i. \quad (12)$$

Example 4.4. Let r and t be positive integers such that $r - 1 \leq \frac{t}{2}$. Then by Lemma 4.2, the 2-orbit scheme of $\mathrm{PGL}(t, q)$ on $\Omega_{r-1} \cup \Omega_{t-r+1}$ is a reduced $(\left[\begin{smallmatrix} t \\ r-1 \end{smallmatrix} \right]_q, 2, r)$ -scheme, say $\mathcal{C}$. Moreover, as the actions of $\mathrm{PGL}(t, q)$ on both Ω_{r-1} and Ω_{t-r+1} are multiplicity free, both $\mathcal{C}_{\Omega_{r-1}}$ and $\mathcal{C}_{\Omega_{t-r+1}}$ are commutative and hence by Corollary 3.1, $|\mathcal{P}(\mathcal{C})| = r$.

Lemma 4.3. *Let $\mathcal{C}_i$ be an (m_i, n_i, r_i) -scheme for $i = 1, 2$. Then $\mathcal{C}_1 \otimes \mathcal{C}_2$ is an $(m_1 m_2, n_1 n_2, r_1 r_2)$ -scheme. Furthermore, $\mathcal{C}_1 \otimes \mathcal{C}_2$ is reduced if and only if both $\mathcal{C}_1$ and $\mathcal{C}_2$ are reduced.*

Proof. The first statement is obtained by the definition of $\mathcal{C}_1 \otimes \mathcal{C}_2$. Let R_i be a basis relation of $\mathcal{C}_i$ for $i = 1, 2$. Then $R_1 \otimes R_2$ is thin if and only if both R_1 and R_2 are thin. This implies that $\mathcal{C}_1 \otimes \mathcal{C}_2$ is reduced if and only if both $\mathcal{C}_1$ and $\mathcal{C}_2$ are reduced. $\square$

Problem 1: Given an odd prime p does there exist any reduced $(m, 3, p)$ -scheme for some m ?

The following problem is inspired from a conjecture by H. Weilandt on permutation representations (see [1], Remark 5.3 and Lemma 5.4).

Problem 2: If $\mathcal{C}$ is a reduced (p, n, r) -scheme for some r and prime p , then $n \leq 2$.

5. Enumeration of (m, n, r) -schemes for $m \leq 11$

Proof of Theorem 1.3 (i): Let $\mathcal{C}$ be a reduced (m, n, r) -scheme and $X, Y \in \mathrm{Fib}(\mathcal{C})$ with $X \neq Y$. Then $2 \leq d_R$ for each $R \in \mathcal{R}_{X,Y}$ and

$$2|\mathcal{R}_{X,Y}| \leq \sum_{R \in \mathcal{R}_{X,Y}} d_R = m,$$

a contradiction.

In order to prove Theorem 1.3 (ii) we need the following lemma.

Lemma 5.1. *Let $\mathcal{C}$ be an (m, n, r) -scheme and $X, Y, Z \in \mathrm{Fib}(\mathcal{C})$. If $T \in \mathcal{R}_{X,Y}$ such that d_T is prime to $\prod_{R \in \mathcal{R}_{Y,Z}} d_R$, then $d_{Y,Z}$ coincides with $d_{X,Z}$ as multi-sets and $d_T \leq \min \{d_R \mid R \in \mathcal{R}_{Y,Z}\}$.*

Proof. For each $R \in \mathcal{R}_{Y,Z}$, $\gcd(d_T, d_R) = 1$. By Lemma 2.2 (viii), $|TR| = 1$ and we may define the following map.

$$\begin{aligned} \psi : \mathcal{R}_{Y,Z} &\longrightarrow \mathcal{R}_{X,Z} \\ R &\mapsto S; \quad TR = \{S\}. \end{aligned}$$

By Lemma 2.2, ψ is surjective. Since $|\mathcal{R}_{Y,Z}| = |\mathcal{R}_{X,Z}|$, ψ must be a bijection. Consequently, $\sum_{R \in \mathcal{R}_{Y,Z}} d_R = \sum_{S \in \mathcal{R}_{X,Z}} d_S = \sum_{R \in \mathcal{R}_{Y,Z}} d_{TR}$. On the other hand, by Lemma 2.2 (v), $d_R \leq d_{TR}$ for each $R \in \mathcal{R}_X$ and thus $d_R = d_{TR}$ for each $R \in \mathcal{R}_{Y,Z}$. Furthermore, by Lemma 2.2 (v), $d_T \leq d_{TR} = d_R$ for each $R \in \mathcal{R}_{Y,Z}$. $\square$

Proof of Theorem 1.3 (ii): Let $\mathcal{C}$ be a reduced (m, n, r) -scheme and let $X \in \text{Fib}(\mathcal{C})$ such that $\mathcal{C}_X$ is p -valenced. Clearly $m = \sum_{T \in \mathcal{R}_{X,Y}} d_T$ where $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$. Since $p \nmid m$, so there exists $T \in \mathcal{R}_{X,Y}$ such that $p \nmid d_T$. Since $\mathcal{C}_X$ is p -valenced, d_T is prime to $\prod_{R \in \mathcal{R}_X} d_R$. As $d_{\Delta_X} = 1$, it follows from Lemma 5.1 that $d_T \leq \min \{d_R \mid R \in \mathcal{R}_X\} = 1$, a contradiction. $\square$

Lemma 5.2. *Let $\mathcal{C}$ be an $(m, n, 2)$ -scheme and $R \in \mathcal{R}_{X,Y}$ where $X, Y \in \text{Fib}(\mathcal{C})$. Then $d_R(d_R - 1) = \lambda(m - 1)$ for some $\lambda \in \mathbb{N}$.*

Proof. Let $\mathcal{C}$ be an $(m, n, 2)$ -scheme and $X, Y \in \text{Fib}(\mathcal{C})$. For each $R \in \mathcal{R}_{X,Y}$ we have by Lemma 2.2 (i),

$$A_R A_{R^t} = \sum_{S \in \mathcal{R}_X} c_{RR^t}^S A_S = d_R I_X + c_{RR^t}^{\Delta_X^c} (J_X - I_X),$$

where $\Delta_X^c = (X \times X) \setminus \Delta_X$. It follows that $R \in \mathcal{R}_{X,Y}$ is regarded as the incident relation of a symmetric (m, d_R, λ) -design where $\lambda = c_{RR^t}^{\Delta_X^c}$. A basic property of symmetric designs implies that $d_R(d_R - 1) = \lambda(m - 1)$. $\square$

Remark 5.3. *Let m and t be positive integers and q an odd prime power such that*

$$m - 1 = 2^t q. \tag{13}$$

Then there are exactly four $d \in \{1, \dots, m - 1\}$ such that $d(d - 1) \equiv 0 \pmod{2^t q}$ by an elementary number theoretical argument. It follows that if $\mathcal{C}$ is a reduced $(m, n, 2)$ -scheme, then $d_{X,Y}$ is uniquely determined for all

$X, Y \in \text{Fib}(\mathcal{C})$. In particular, if m is prime satisfying (13), then there is $\gamma \in \{2, \dots, m-2\}$ such that $d_{X,Y} = \{\gamma, m-\gamma\}$ and $\gcd(\gamma, m-\gamma) = 1$ for all $X, Y \in \text{Fib}(\mathcal{C})$.

Lemma 5.4. *Let $\mathcal{C}$ be a reduced $(m, n, 2)$ -scheme. Suppose that $d_{X,Y} = \{a, b\}$ with $\gcd(a, b) = 1$ for all $X, Y \in \text{Fib}(\mathcal{C})$. Then $n \leq 2$.*

Proof. Suppose that X, Y and Z are distinct fibers of $\mathcal{C}$ and let $\mathcal{R}_{X,Y} = \{R, R'\}$, $\mathcal{R}_{Y,Z} = \{S, S'\}$, $\mathcal{R}_{X,Z} = \{T, T'\}$ so that $d_R = d_S = d_T = a < b = d_{R'} = d_{S'} = d_{T'}$. By Lemma 2.2 (i), (ii), (iii), $a^2 = d_R d_S = \alpha a + \beta b$ such that $a \mid b\beta$ and $\beta < a$. Since $\gcd(a, b) = 1$, it follows that $\beta = 0$ and $\alpha = a$. This implies that $c_{RS}^T = a = d_R$. It follows that

$$R_{\text{out}}(x) \subseteq S_{\text{in}}(z), \quad (14)$$

where $(x, z) \in T$. Now we take $y_1, y_2 \in R_{\text{out}}(x)$ so that $y_1 \neq y_2$. It follows from (14) that $T_{\text{out}}(x) \subseteq S_{\text{out}}(y_1) \cap S_{\text{out}}(y_2)$. This fact along with Lemma 2.2 (iii) assert that $a = c_{SSt}^{\Delta_Y^c}$ where $\Delta_Y^c = (Y \times Y) \setminus \Delta_Y$. Therefore, by Lemma 5.2, $a(a-1) = a(a+b-1)$. It follows that $ab = 0$, a contradiction. This completes the proof. $\square$

Lemma 5.5. *Let $\mathcal{C}$ be a reduced $(m, n, 2)$ -scheme. If $m-1$ is a prime power, then $n = 1$.*

Proof. Let p be prime such that $m-1 = p^t$ for some t . In this case, p does not divide m and we are done by Theorem 1.3 (ii). $\square$

Lemma 5.6. *Let $\mathcal{C}$ be a reduced (m, n, r) -scheme and $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$. If $m = 2r$, then the following hold:*

- (i) *For each $T \in \mathcal{R}_{X,Y}$, $d_T = 2$.*
- (ii) *For each $R \in \mathcal{R}_X$, $d_R \in \{1, 2, 4\}$ and*

$$|\{R \in \mathcal{R}_X \mid d_R = 1\}| = 2|\{R \in \mathcal{R}_X \mid d_R = 4\}|.$$

Proof. (i) Let $\mathcal{C}$ be a reduced (m, n, r) -scheme and $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$. Then as $d_T \geq 2$ for each $T \in \mathcal{R}_{X,Y}$, it follows from $m = \sum_{T \in \mathcal{R}_{X,Y}} d_T$ that $2r \leq m$ and the equality holds if and only if $d_T = 2$ for each $T \in \mathcal{R}_{X,Y}$.

(ii) Let $R \in \mathcal{R}_X$ and $T \in \mathcal{R}_{X,Y}$. Then by Lemma 2.2 (i), (iii), there exist none-negative integers α and β such that $2d_R = d_R d_T = \alpha d_S + \beta d_{S'} = 2\alpha + 2\beta$ and $\alpha, \beta \leq 2$. This implies that $d_R \leq 4$. By Lemma 5.1, $d_R \in \{1, 2, 4\}$. We set $k_i := |\{R \in \mathcal{R}_X \mid d_R = i\}|$ for $i \in \{1, 2, 4\}$. Since $k_1 + k_2 + k_4 = |\mathcal{R}_X| = |\mathcal{R}_{X,Y}|$, it follows that $m = k_1 + 2k_2 + 4k_4 = 2(k_1 + k_2 + k_4)$. Therefore, $k_1 = 2k_4$. $\square$

Lemma 5.7. *For each (m, n, r) -scheme, if m is prime, then $r - 1$ divides $m - 1$.*

Proof. Let $X \in \text{Fib}(\mathcal{C})$ and consider the homogeneous component $(X, \mathcal{R}_X)$. Since $|X| = m$ is prime, by [7, Theorem 3.3] $d_R = d$ for all $R \in \mathcal{R}_X$ with $R \neq \Delta_X$. Then $m - 1 = \sum_{\substack{R \in \mathcal{R}_X \\ \Delta_X \neq R}} d_R = (r - 1)d$. $\square$

Lemma 5.8. *Let $\mathcal{C} = (V, \mathcal{R})$ be an (m, n, r) -scheme. If m is odd, then each non-reflexive symmetric basis relation of $\mathcal{C}$ has even degree.*

Proof. Let $S \in \mathcal{R}_X$ be symmetric for some $X \in \text{Fib}(\mathcal{C})$. Since $S \neq \Delta_X$, $|S|$ is even. By (2), $|S| = d_S m$ and thus d_S is even. $\square$

Lemma 5.9 ([10, (3.2)]). *Let $\mathcal{C}$ be a reduced $(m, n, 3)$ -scheme. Then $\mathcal{C}_X$ is symmetric for each $X \in \text{Fib}(\mathcal{C})$.*

Proof of Theorem 1.4: So far in this section we have been preparing some lemmas which will be applied to enumerate reduced (m, n, r) -schemes for m up to 11.

The entries (r, m) such that $m < 2r$ are eliminated by Theorem 1.3 (i) whereas $(2, m)$'s are eliminated by Lemma 5.5 except $(2, 7)$ and $(2, 11)$. If $\mathcal{C}$ is a reduced $(m, n, 2)$ -scheme with $m \in \{7, 11\}$, then by Remark 5.3 and Lemma 5.4, $n \leq 2$. Thus we have eliminated the first row of Table 1.

Applying Lemma 5.6 for $(r, m) = (5, 10)$ we obtain that $d_X = \{1, 1, 2, 2, 4\}$. According to [6, 13] there are no homogeneous schemes with $d_X = \{1, 1, 2, 2, 4\}$ where we can prove this fact by a theoretical way.

The entries $(4, 11)$ and $(5, 11)$ are eliminated by Lemma 5.7 whereas $(3, 7)$ and $(3, 11)$ are eliminated by Lemmas 5.9 and 5.8.

An (r, m) -entry of the following table is denoted by $*$ if there exists no $(m, 1, r)$ -scheme.

$r \backslash m$	4	5	6	7	8	9	10	11
2	1	1	1	≤ 2	1	1	1	≤ 2
3	1	1	1	1	≤ 2	1	1	1
4	1	*	1	1	≤ 2	1	1	*
5	*	1	*	*	1	1	*	*

Table 1.

The following is the list of (r, m) , $\sum_{i=1}^r a_i$ and $\sum_{i=1}^r b_i$ where $m = \sum_{i=1}^r a_i = \sum_{i=1}^r b_i$, $1 = a_1 \leq a_2 \leq \dots \leq a_r$ and $2 \leq b_1 \leq b_2 \leq \dots \leq b_r$ such that $d_X = \{a_1, \dots, a_r\}$ and $d_{X,Y} = \{b_1, \dots, b_r\}$ for some $(m, 1, r)$ -scheme $(X, \mathcal{R}_X)$ not satisfying the assumption of Theorem 1.3 (see [6, 13]).

(r, m)	$\sum_{R \in \mathcal{R}_X} d_R$	$\sum_{R \in \mathcal{R}_{X,Y}} d_R$	
(3,6)	1+1+4	2+2+2	Not occur by Lemma 5.10
(3,8)	1+1+6	2+2+4	Not occur by Lemma 5.11
		2+3+3	$n \leq 2$ by Lemma 5.12
	1+3+4	2+2+4	Not occur by Lemma 2.2 (vii)
		2+3+3	Not occur by Lemma 2.2 (vii)
(3,9)	1+2+6	2+2+5	Not occur by Lemma 5.1
		2+3+4	Not occur by Lemma 5.13
		3+3+3	Not occur (see Lemma 5.13)
(3,10)	1+1+8	2+3+5	Not occur by Lemma 5.1
		3+3+4	Not occur by Lemma 5.1
		2+4+4	Not occur by Lemma 5.14 (i)
		2+2+6	Not occur by Lemma 5.14 (i)
	1+3+6	2+3+5	Not occur by Lemma 5.1
		3+3+4	Not occur by Lemma 5.14 (ii)
		2+4+4	Not occur by Lemma 5.1
		2+2+6	Not occur by Lemma 2.2 (vii)
	1+4+5	2+3+5	Not occur by Lemma 5.1
		3+3+4	Not occur by Lemma 5.1
		2+4+4	Not occur by Lemma 2.2 (vii)
		2+2+6	Not occur by Lemma 2.2 (vii)
(4,8)	1+1+2+4	2+2+2+2	$n \leq 2$ (see Lemma 5.16)
(4,9)	1+1+1+6	2+2+2+3	Not occur by Lemma 2.2 (vii) and Lemma 5.8
	1+2+3+3	2+2+2+3	Not occur by Lemma 5.17
(4,10)	1+2+2+5	2+2+2+4	Not occur by Lemma 5.1
		2+2+3+3	Not occur by Lemma 5.1
	1+1+4+4	2+2+3+3	Not occur by Lemma 5.1
		2+2+2+4	Not occur by Lemma 5.18

Table 2.

Lemma 5.10. *If $\mathcal{C}$ is a reduced $(6, n, 3)$ -scheme such that $d_X = \{1, 1, 4\}$ for some $X \in \text{Fib}(\mathcal{C})$, then $d_{X,Y} \neq \{2, 2, 2\}$ for each $Y \in \text{Fib}(\mathcal{C})$ with $Y \neq X$.*

Proof. Suppose by the contrary that $d_{X,Y} = \{2, 2, 2\}$ for some $Y \neq X$. By Lemma 5.6, $d_Y = \{1, 1, 4\}$. Taking $R, S \in \mathcal{R}_{X,Y}$ with $R \neq S$ we obtain from Lemma 2.2 (vii) that $R^t R = S^t S = \{\Delta_Y, T\}$ where $T \in \mathcal{R}_Y$ with $T \neq \Delta_Y$ and $d_T = 1$. By Lemma 2.2 (i), (ii), (iv), $4 = d_R d_{S^t} = \alpha + 4\beta$ for some non-negative integers $\alpha, \beta \leq 2$. This implies $\alpha = 0$ and $\beta = 1$, which contradicts Lemma 2.4. $\square$

Lemma 5.11. *If $\mathcal{C}$ is a reduced $(8, n, 3)$ -scheme such that $d_X = \{1, 1, 6\}$ for some $X \in \text{Fib}(\mathcal{C})$, then we have the following:*

- (i) *For each $Y \in \text{Fib}(\mathcal{C})$ with $Y \neq X$, $d_{X,Y} \neq \{2, 2, 4\}$. Indeed, $d_{X,Y} = \{2, 3, 3\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*
- (ii) *Let $\mathcal{R}_{X,Y} = \{R, S, S'\}$ such that $d_R = 2$ and $d_S = d_{S'} = 3$. Let $T \in \mathcal{R}_X$ with $T \neq \Delta_X$ and $d_T = 1$. Then $TR = \{R\}$, $TS = \{S'\}$ and $TS' = \{S\}$.*

Proof. (i) Suppose by the contrary that $d_{X,Y} = \{2, 2, 4\}$ for some $Y \in \text{Fib}(\mathcal{C})$, and take $R \in \mathcal{R}_X$ and $S \in \mathcal{R}_{X,Y}$ so that $d_R = 6$ and $d_S = 2$. It follows from Lemma 2.2 (i), (ii), (iii) that for some non-negative integers α, β, γ we have

$$12 = d_R d_S = 2\alpha + 2\beta + 4\gamma, \quad 6 \mid 2\alpha, \quad 6 \mid 2\beta, \quad 6 \mid 4\gamma \quad \alpha, \beta, \gamma \leq 2.$$

This implies that $\gamma = 0$ and $12 = 2\alpha + 2\beta \leq 8$, a contradiction.

(ii) As $d_T = 1$, Lemma 2.2 (v), (viii) asserts that $d_{TR} = 2$ and $TR = \{R\}$, since R is the unique basis relation in $\mathcal{R}_{X,Y}$ of degree 2. By the same observation $d_{TS} = 3$ and $TS \in \mathcal{R}_{X,Y}$. If $TS = \{S\}$, then by Lemma 2.2 (i), $c_{TS}^S = 1$ and Lemma 2.2 (ii) implies that $c_{SS^t}^T = 3$. Therefore, applying Lemma 2.2 (i), (iv) we have $9 = d_S d_{S^t} = 3 + 3 + c_{SS^t}^{T'} 6$ where $T' \in \mathcal{R}_X$ with $d_{T'} = 6$, a contradiction. $\square$

Lemma 5.12. *If $\mathcal{C}$ is a reduced $(8, n, 3)$ -scheme such that $d_X = \{1, 1, 6\}$ for some $X \in \text{Fib}(\mathcal{C})$, then $n \leq 2$.*

Proof. Suppose by the contrary that X, Y and Z are distinct fibers of $\mathcal{C}$. Then by Lemma 5.11, $d_{X,Y} = d_{Y,Z} = d_{X,Z} = \{2, 3, 3\}$. Let $R \in \mathcal{R}_{X,Y}$ and

$S \in \mathcal{R}_{Y,Z}$ with $d_R = 2$ and $d_S = 3$. It follows from Lemma 2.2 (i), (ii), (iii), there exist non-negative integers α, β, γ such that

$$6 = d_R d_S = 2\alpha + 3\beta + 3\gamma, \quad 6 \mid 2\alpha, \quad \alpha \leq 2.$$

This implies $\alpha = 0$ and $RS = \{S'\}$ where $S' \in \mathcal{R}_{Y,Z}$ with $d_{S'} = 3$. Since $c_{RS}^{S'} = 2$, by Lemma 2.4, $R^t R \cap SS^t = \{\Delta_Y, T\}$ for some $T \in \mathcal{R}_Y$ with $d_T = 1$. Thus $9 = d_S d_{S'} = 3 + 3 + 6\alpha$. It follows that $3 = 6\alpha$, a contradiction. $\square$

Lemma 5.13. *Let $\mathcal{C}$ be a reduced $(9, n, 3)$ -scheme and $d_X = \{1, 2, 6\}$ for some fiber X . Then $d_{X,Y} \notin \{\{2, 3, 4\}, \{3, 3, 3\}\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*

Proof. Suppose that $d_{X,Y} = \{2, 3, 4\}$ and take the basis relations $R \in \mathcal{R}_X$ and $S \in \mathcal{R}_{X,Y}$ so that $d_R = 6$ and $d_S = 2$. It follows from Lemma 2.2 (i), (ii), (iii), there exist non-negative integers α, β, γ such that

$$12 = d_R d_S = 2\alpha + 3\beta + 4\gamma, \quad 6 \mid 2\alpha, \quad 6 \mid 3\beta, \quad 6 \mid 4\gamma, \quad \alpha, \beta, \gamma \leq 2.$$

This implies that $\gamma = 0$ and $12 = 2\alpha + 3\beta \leq 10$, a contradiction.

Suppose that $d_{X,Y} = \{3, 3, 3\}$ for some $Y \in \text{Fib}(\mathcal{C})$ and take distinct $R, S \in \mathcal{R}_{X,Y}$. By Lemma 2.2 (i), (iv), for some non-negative integers α, β we have $9 = d_R d_{S^t} = 2\alpha + 6\beta = 2(\alpha + 3\beta)$, a contradiction. $\square$

Let $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$ and $R, S, S' \in \mathcal{R}_{X,Y}$. Then $R^t R \cap S^t S' \neq \emptyset$ if and only if $RS^t \cap RS'^t \neq \emptyset$. We use this fact in the proof of the following lemma.

Lemma 5.14. *Let $\mathcal{C}$ be a reduced $(10, n, 3)$ -scheme. Then the following hold:*

- (i) *If $d_X = \{1, 1, 8\}$ for some $X \in \text{Fib}(\mathcal{C})$, then $d_{X,Y} \notin \{\{2, 2, 6\}, \{2, 4, 4\}\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*
- (ii) *If $d_X = \{1, 3, 6\}$ for some $X \in \text{Fib}(\mathcal{C})$, then $d_{X,Y} \neq \{3, 3, 4\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*

Proof. (i) Suppose that $d_{X,Y} = \{2, 2, 6\}$ for some $Y \in \text{Fib}(\mathcal{C})$. Take $R, S \in \mathcal{R}_{X,Y}$ with $R \neq S$ and $d_R = d_S = 2$. By Lemma 2.2 (i), (iv), (iii), $4 = d_R d_{S^t} = \alpha + 8\beta$ for some non-negative integers α, β with $\alpha, \beta \leq 2$. It follows that $\alpha = 0$ and $4 = 8\beta$, a contradiction.

Suppose that $\mathcal{R}_{X,Y} = \{R, S, S'\}$ such that $d_R = 2$ and $d_S = d_{S'} = 4$. By Lemma (i), (iv), $8 = d_R d_{S^t} = \alpha + 8\beta$ for some non-negative integers α, β with

$4 \mid \alpha \leq 2$. This implies that $\alpha = 0$ and then $\beta = 1$. Therefore, $RS^t = \{T'\}$ where $T' \in \mathcal{R}_X$ with $d_{T'} = 8$. By the same observation, $RS^{t'} = \{T'\}$. Therefore, $T \in S^t S' \cap R^t R$ where $T \neq \Delta_Y$. On the other hand, by Lemma 2.2 (vii), $d_T = 1$. It follows from Lemma 2.2 (i), (ii), (iv), (iii) that for some non-negative integers α, β we have $16 = d_{S^t} d_{S'} = \alpha + 8\beta$ with $4 \mid \alpha$ and $0 < \alpha \leq 4$. This implies that $\alpha = 4$ and thus $12 = 8\beta$, a contradiction.

(ii) Take $R \in \mathcal{R}_X$ and $S \in \mathcal{R}_{X,Y}$ with $d_R = 3$ and $d_S = 4$. By Lemma 2.2 (i), (ii), (iii), for some non-negative integers α, β, γ we have $12 = d_R d_S = 3\alpha + 3\beta + 4\gamma$ with $12 \mid \alpha$ and $12 \mid \beta$ and $\alpha, \beta, \gamma \leq 3$. This implies that $\alpha = \beta = 0$ and $\gamma = 3$. Hence $RS = \{S\}$. By Lemma 2.3, $d_{L_S} \mid \gcd(10, 4) = 2$ which is a contradiction since $d_{L_S} > d_R = 3$. $\square$

Lemma 5.15. *Let $\mathcal{C}$ be a reduced $(8, n, 4)$ -scheme such that $d_X = \{1, 1, 2, 4\}$ for some $X \in \text{Fib}(\mathcal{C})$. Then for all $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$, there exists $R \in \mathcal{R}_{X,Y}$ such that $RR^t = \{\Delta_X, S\}$ (resp. $R^t R = \{\Delta_Y, S'\}$) where S is the unique basis relation in $\mathcal{R}_X$ with $d_S = 2$ (resp. S' is the unique basis relation in $\mathcal{R}_Y$ with $d_{S'} = 2$).*

Proof. Let $\mathcal{C}$ be a reduced $(8, n, 4)$ -scheme such that $d_X = \{1, 1, 2, 4\}$ for some $X \in \text{Fib}(\mathcal{C})$. Then $d_{X,Y} = \{2, 2, 2, 2\}$ for all $X, Y \in \text{Fib}(\mathcal{C})$ with $X \neq Y$. Let $T \in \mathcal{R}_X$ with $T \neq \Delta_X$ and $d_T = 1$. Then $d_{TR} = 2$ and $|TR| = 1$. Suppose that $TR = \{R\}$ for each $R \in \mathcal{R}_{X,Y}$. Then $T \notin RS^t$ for all $R, S \in \mathcal{R}_{X,Y}$ with $R \neq S$. Thus by Lemma 2.2 (i), (iii), $4 = d_R d_{S^t} = 2\alpha + 4\beta$ for some non-negative integers α, β with $\alpha \leq 2$. By Lemma 2.4, $\beta = 0$ and $\alpha = 2$. This implies that $R\mathcal{R}_{Y,X} \subsetneq \mathcal{R}_X$, which contradicts Lemma 2.2 (iii). Thus there exists $R \in \mathcal{R}_{X,Y}$ such that $TR \neq \{R\}$. Equivalently, $T \notin RR^t$. It follows from Lemma 2.2 (vii) that $RR^t = \{\Delta_X, S\}$ where S is the unique basis relation in $\mathcal{R}_X$ with $d_S = 2$. $\square$

Lemma 5.16. *If $\mathcal{C}$ is a reduced $(8, n, 4)$ -scheme such that $d_X = \{1, 1, 2, 4\}$ for some $X \in \text{Fib}(\mathcal{C})$, then $n \leq 2$.*

Proof. Suppose by the contrary that X, Y and Z are distinct fibers of $\mathcal{C}$. Then by Lemma 5.15, there exist $R \in \mathcal{R}_{X,Y}$ and $T \in \mathcal{R}_{Y,Z}$ such that $R^t R = TT^t = \{\Delta_X, S\}$ where S is the unique basis relation in $\mathcal{R}_Y$ with $d_S = 2$. It follows from Lemma 2.2 (i) that $c_{TT^t}^S = c_{R^t R}^S = 1$. Let $(y, y') \in S$. Then there exists $(x, z) \in X \times Z$ such that $R_{in}(y) \cap R_{in}(y') = \{x\}$ and $T_{out}(y) \cap T_{out}(y') = \{z\}$. As $d_T = 2$, we may assume that $T_{out}(y) = \{z, z_1\}$ and $T_{out}(y') = \{z, z_2\}$. Note that $z_1 \neq z_2$, otherwise $c_{TT^t}^S \geq 2$, a contradiction. This means that

$(R \circ T)_{out}(x) = \{z, z_1, z_2\}$ and thus $d_{RT} = d_{R \circ T} = 3$, which is a contradiction since d_{RT} must be a sum of degrees in $d_{X,Z} = \{2, 2, 2, 2\}$.




Figure 2.

□

Example 5.1. The association scheme **as16** No.122 as in [6] induces the thin residue fission (see [12, Proposition 3.1]), which is a reduced $(8, 2, 4)$ -scheme whose relational matrix is

$$\left(\begin{array}{cccccccc|cccccccc} 0 & 1 & 2 & 2 & 3 & 3 & 3 & 3 & 4 & 4 & 5 & 5 & 6 & 7 & 6 & 7 \\ 1 & 0 & 2 & 2 & 3 & 3 & 3 & 3 & 4 & 4 & 5 & 5 & 7 & 6 & 7 & 6 \\ 2 & 2 & 0 & 1 & 3 & 3 & 3 & 3 & 5 & 5 & 4 & 4 & 6 & 7 & 7 & 6 \\ 2 & 2 & 1 & 0 & 3 & 3 & 3 & 3 & 5 & 5 & 4 & 4 & 7 & 6 & 6 & 7 \\ 3 & 3 & 3 & 3 & 0 & 1 & 2 & 2 & 6 & 7 & 6 & 7 & 4 & 4 & 5 & 5 \\ 3 & 3 & 3 & 3 & 1 & 0 & 2 & 2 & 7 & 6 & 7 & 6 & 4 & 4 & 5 & 5 \\ 3 & 3 & 3 & 3 & 2 & 2 & 0 & 1 & 6 & 7 & 7 & 6 & 5 & 5 & 4 & 4 \\ 3 & 3 & 3 & 3 & 2 & 2 & 1 & 0 & 7 & 6 & 6 & 7 & 5 & 5 & 4 & 4 \\ \hline 4' & 4' & 5' & 5' & 6' & 7' & 6' & 7' & 0' & 1' & 2' & 2' & 3' & 3' & 3' & 3' \\ 4' & 4' & 5' & 5' & 7' & 6' & 7' & 6' & 1' & 0' & 2' & 2' & 3' & 3' & 3' & 3' \\ 5' & 5' & 4' & 4' & 6' & 7' & 7' & 6' & 2' & 2' & 0' & 1' & 3' & 3' & 3' & 3' \\ 5' & 5' & 4' & 4' & 7' & 6' & 6' & 7' & 2' & 2' & 1' & 0' & 3' & 3' & 3' & 3' \\ 6' & 7' & 6' & 7' & 4' & 4' & 5' & 5' & 3' & 3' & 3' & 3' & 0' & 1' & 2' & 2' \\ 7' & 6' & 7' & 6' & 4' & 4' & 5' & 5' & 3' & 3' & 3' & 3' & 1' & 0' & 2' & 2' \\ 6' & 7' & 7' & 6' & 5' & 5' & 4' & 4' & 3' & 3' & 3' & 3' & 2' & 2' & 0' & 1' \\ 7'' & 6' & 6' & 7'' & 5' & 5' & 4' & 4' & 3' & 3' & 3' & 3' & 2' & 2' & 1' & 0' \end{array} \right).$$

Also the thin residue fission of the association scheme **as16** No.51 as in [6], is a reduced $(8, 2, 3)$ -scheme.

Lemma 5.17. *Let $\mathcal{C}$ be a reduced $(9, n, 4)$ -scheme such that $d_X = \{1, 2, 3, 3\}$ for some $X \in \text{Fib}(\mathcal{C})$. Then $d_{X,Y} \neq \{2, 2, 2, 3\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*

Proof. Suppose by the contrary that $R_1, R_2, R_3 \in \mathcal{R}_{X,Y}$ with $d_{R_i} = 2$ for $i \in \{1, 2, 3\}$. For all $i, j \in \{1, 2, 3\}$ with $i \neq j$, by Lemma 2.2 (i), (iv) we have $4 = d_{R_i}d_{R_j^t} = 2\alpha + 3\beta + 3\gamma$. This implies that $\beta = \gamma = 0$ and $\alpha = 2$. Hence, for all $i, j \in \{1, 2, 3\}$ with $i \neq j$, $R_iR_j^t = \{T\}$ where $T \in \mathcal{R}_X$ with $d_T = 2$. It follows from Lemma 2.2 (vi) that $\{R_1\} = TR_2 = \{R_3\}$, a contradiction. $\square$

Lemma 5.18. *Let $\mathcal{C}$ be a reduced $(10, n, 4)$ -scheme such that $d_X = \{1, 1, 4, 4\}$ for some $X \in \text{Fib}(\mathcal{C})$. Then $d_{X,Y} \neq \{2, 2, 2, 4\}$ for each $Y \in \text{Fib}(\mathcal{C})$.*

Proof. Suppose by the contrary that $d_{X,Y} = \{2, 2, 2, 4\}$ for some $Y \in \text{Fib}(\mathcal{C})$ with $Y \neq X$. According to [6, 13], $d_Y \in \{\{1, 1, 4, 4\}, \{1, 2, 2, 5\}\}$. It follows from Lemma 5.1 that $d_Y = \{1, 1, 4, 4\}$. Take $R, S \in \mathcal{R}_{X,Y}$ with $R \neq S$ and $d_R = d_S = 2$. By Lemma 2.2 (i), (iv), (iii), there exist non-negative integers α, β, γ such that

$$4 = d_Rd_{S^t} = \alpha + 4\beta + 4\gamma, \quad \alpha, \beta, \gamma \leq 2.$$

This implies that $4 \mid \alpha$. Hence, $\alpha = 0$ and $\beta + \gamma = 1$ which contradicts Lemma 2.4. $\square$

Acknowledgements

The authors would like to thank I. Ponomarenko, E. Bannai and A. Ivanov for their valuable comments to this article, especially internal direct sums, $(m, 2, r)$ -schemes and tensor products, respectively.

- [1] P. J. Cameron, “ *On groups with several doubly-transitive permutation representations*”, Math. Z. 128(1972), 1-14.
- [2] P. J. Cameron, “ *Combinatorics and Groups: Peter Cameron’s IPM Lecture Notes*”, Published by IPM, Tehran, 2001.
- [3] S. Evdokimov and I. Ponomarenko, “ *Permutation group approach to association schemes*”, European Journal of Combinatorics, 30(2009), 1456-1476.
- [4] S. Evdokimov and I. Ponomarenko, “ *Two inequalities for the parameters of a coherent algebra*”, Zap. Nauchnykh Seminarov POMI 240(1997), 82-95. English translation: J. Math. Sci., New York 96(5)(1999), 3496-3504.
- [5] A. Hanaki, “ *Representations of finite association schemes*”, European Journal of Combinatorics, 30(2009), 1477-1496.

- [6] A. Hanaki and I. Miyamoto, “ *Classification of association schemes of small order, Online catalogue*”, available from <http://kissme.shinshu-u.ac.jp/as>
- [7] A. Hanaki, K. Uno, “ *Algebraic structure of association schemes of prime order*”, J. Algebraic. Combin. 23 (2)(2006), 189 - 195.
- [8] D. G. Higman, “ *Coherent configuration part I: Ordinary representation theory*”, Geometriae Dedicata (1975), 1-32.
- [9] D. G. Higman, “ *Coherent algebras*”, Linear Algebra Appl., 93(1987), 209-239.
- [10] D. G. Higman, “ *Strongly regular designs of the second kind*”, European Journal of Combinatorics, 16(1995), 479-490.
- [11] M. Hirasaka, “ *The classification of association schemes with 11 and 12 vertices*”, Kyushu J. Math. 51(1997) 413-428.
- [12] M. Klin, M. Muzychuk, C. Pech, A. Woldar, and P.-H. Zieschang, “ *Association schemes on 28 points as mergings of a half-homogeneous coherent configuration*”, European Journal of Combinatorics, (28)2007, 994-2025.
- [13] E. Nomiyama, “ *Classification of associatian schemes with at most ten vertices*”, Kyushu J. Math. 49(1995) 163-195.
- [14] H. Pollatsek, “ *First cohomology groups of some linear groups over fields of characteristic two*”, Illinois J. Math. 15(1971), 393-417.
- [15] P.-H. Zieschang, “ *Theory of association schemes*”, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2005.